

Dhofar 1180

Polymict regolith breccia

115 g



Figure 1: Dhofar 1180 illustrating dark glassy matrix and light colored feldspathic clasts. Image from M. Farmer. Width of sample is 8 cm.

Introduction

Dhofar 1180 (Fig. 1) was found in the Dhofar region of Oman in January 2005 (Figs. 2 and 3). The meteorite has an external shape similar to a "thick-bladed talon" (Connolly et al., in preparation). The sample is a polymict regolith breccia that contains a variety of lithologies set in a matrix of similar materials with a preferred orientation of fragments and clasts (Fig. 4).




Figure 2 and 3: Location maps of the Dhofar region in Oman (from Al-Kathiri et al., 2005) and the specific coordinates for Dhofar 1180 (near center).

Petrography and mineralogy

Clasts (0.1 to 1 mm) in this meteorite are mostly ferroan anorthosites (Fa_{38} ; $\text{Fs}_{38.6}\text{Wo}_{2.1}$) as well as gabbroic anorthosites, anorthositic gabbros, norites (Fa_{18}), troctolites, olivine gabbros ($\text{Fa}_{36.8}$; $\text{Fs}_{33.4}\text{Wo}_{4.3}$), microporphyritic and fine-grained impact melt breccias, and rare, ophitic/subophitic basalts (pyroxene core - $\text{Fs}_{40}\text{Wo}_{11.8}$, rim - $\text{Fs}_{69.1}\text{Wo}_{15.8}$) (Connolly et al., 2006; Bunch et al., 2006; Zhang and Hsu, 2006, 2007, 2009). Zhang and Hsu (2007, 2009) describe one KREEP related clast, although this component must be minor because the bulk chemistry of Dho1180 is not particularly KREEPy (see below). The plagioclase feldspar varies in composition from An_{91-99} , and the matrix contains numerous fragments of plagioclase, pyroxene, and olivine (Fig. 4 and 5). In addition to the wide variety of clasts (Zhang and Hsu, 2009), there are also a number of glassy fragments in Dho 1180 (Fig. 6), making it a regolith breccias as opposed to simply a fragmental breccias (Zhang and Hsu, 2009).



Figure 4: Cut slab face of Dhofar 1180 illustrating dark glassy matrix and light colored feldspathic clast. Image from M. Farmer. Width of sample is 10 cm.




Figure 5: Thin section image of Dhofar 1180. Image from T. Bunch and T. Irving. Field of view is 5 mm.




Figure 6: BSE images of glasses in Dhofar 1180. a) A mafic glass including some anhedral-subhedral pigeonite grains. b) A mafic glass with a near-circular outline. This glass includes a relict augite grain and plagioclase glass. The glass portion shows devitrification. Some very finegrained metal grains exist in this glass. c) A mafic glass that is attached to a pyroxene grain. d) A mafic glass containing some plagioclase grains, and fragments of exsolved enstatite and K-rich feldspar. Some very fine-grained metal grains exist in this glass. e) A feldspathic glass showing devitrification at the margin. Arrows indicate glass regions. pgt=pigeonite; aug=augite; ol=olivine; pl=plagioclase; kfs=K-rich feldspar; en=enstatite; mt=metal. [from Zhang and Hsu, 2009].

Chemistry

Preliminary compositional data for this meteorite, showed it to have 22.6 wt% Al_2O_3 , 9.3 wt% FeO, 0.9 ppm Th (Bunch et al., 2006), 26.8 ppm Sc, 1040 ppm Cr, 130 ppm Ni, and 2.84 ppm Eu (Korotev et al., 2008). These traits alone demonstrate that it is a mixed breccia that includes both basaltic and highlands materials, but no KREEP. Additional studies have sharpened the understanding of the unique nature of this meteorite. It has lower REE contents than many polymict breccias (Fig. 7). In addition, it overlaps somewhat, but is nonetheless distinct from other polymict lunar meteorite breccias in alkali, Sc and Sm contents, such as the YQN samples and NWA 5153 and 5207 (Fig. 8).




Figure 8: Comparison of Dho 1180 bulk composition to other polymict lunar meteorite breccias, illustrating its distinct composition with slightly higher Eu, lower Sc compared to the YQN pairing group (Yamato, QUE, NWA 2995 pairs; Korotev et al., 2009b).

Table 1a:Chemical composition of Dho 1180

<i>reference</i>	1	1	2	Rh		
<i>weight</i>	20-60	196	91.19	Pd ppb		
<i>technique</i>	a	c	c	Ag ppb		
				Cd ppb		
SiO ₂ %	45.4			In ppb		
TiO ₂	0.71			Sn ppb		
Al ₂ O ₃	22.6			Sb ppb		
FeO	9.22			Te ppb		
MnO	0.15			Cs ppm	<0.2	0.052
MgO	6.17			Ba	537	594
CaO	14.9			La	5.3	4.18
Na ₂ O	0.36			Ce	14.1	10.67
K ₂ O	0.06			Pr		
P ₂ O ₅	0.05			Nd	8.9	6.7
S %				Sm	2.84	2.14
sum	99.7			Eu	0.9	0.856
				Gd		
Sc ppm		26.8	15.53	Tb	0.64	0.453
V				Dy		
Cr		1040	869	Ho		
Co		18.6	15.68	Er		
Ni		128	112	Tm		
Cu				Yb	2.44	1.711
Zn				Lu	0.345	0.239
Ga				Hf	2.18	1.55
Ge				Ta	0.3	0.213
As		0.35		W ppb		
Se		n.a.		Re ppb		
Rb		<8		Os ppb		
Sr		1580	1973	Ir ppb	5.3	3.6
Y				Pt ppb		
Zr		75	47	Au ppb	1.5	10.1
Nb				Th ppm	0.9	0.696
Mo				U ppm	0.32	0.37
Ru						

technique (a) EMPA, (b) ICP-MS, (c) INAA (d) XRF

Table 1b. Light and/or volatile elements for Dho 1180

Li ppm

Be

C

S

F ppm

Cl

Br 0.53

I

Pb ppm

Hg ppb

Tl

Bi

References: 1) Korotev et al. (2009b); 2) Zhang and Hsu (2009)

Radiometric age dating

There are no known studies.

Cosmogenic exposure ages

There are no known studies.

K. Righter – Lunar Meteorite Compendium - 2010