


**LEW88516** – 13.2 grams  
Lherzolitic Shergottite



*Figure 1. Photograph of LEW88516 after first break for processing. Cube is 1 cm for scale. NASA # S91-37060*

### **Introduction**

Lewis Cliff sample LEW88516 is petrologically similar to ALH77005 (Satterwhite and Mason 1991; Treiman *et al.* 1994) and Y793605 (Kojima *et al.* 1997). However, it is not paired, because it has a distinctly different terrestrial exposure age. Figure 1 shows LEW88516 as a small (2 cm), dark, rounded, meteorite with coarsely crystalline interior.

LEW88516 was a small nondescript meteorite that was not recognized as an achondrite until it was broken open. Hence, it waited 2 years to be processed in numerical order. According to Satterwhite and Mason (1991), LEW88516 had a “*pitted and mostly shiny fusion crust over 80 % of the surface.*” Preliminary examination of the thin section showed about 50 % olivine, 35 % pyroxene, 8 % interstitial maskelynite with about 5% brown glass. Grain size is about 2 - 3 mm.

### **Petrography**

Harvey *et al.* (1993), Treiman *et al.* (1994), Gleason *et al.* (1997) and Mikouchi and Miyamoto (2000) have given detailed petrologic descriptions of LEW88516 and report that it is similar to ALHA77005 (figure 2). This


meteorite has three distinct textural regions: 1) poikilitic crystalline, 2) interstitial crystalline (or non-poikilitic) and, 3) glassy to partially crystalline veinlets. Delaney (1992), Lindstrom *et al.* (1992) and Harvey and McSween (1992) have also reported petrological descriptions of LEW88516. Papike *et al.* (2009) compared LEW88516 with the roest of the shergottites.

In the poikilitic regions, mm-sized olivine crystals and 50 micron-sized chromite crystals are contained within, and completely surrounded by, pigeonite crystals with exsolution lamellae of augite. Maskelynite, whitlockite and sulfides are rare in the poikilitic regions.

The interstitial areas have a more typically basaltic texture with intergrown euhedral and subhedral olivine, pigeonite and chromite, with interstitial maskelynite, pigeonite, whitlockite, ilmenite and pyrrhotite.

The glass veinlets are the result of shock melting of the other lithologies (Harvey *et al.* 1993).

Melt inclusions in large olivine grains were studied by



**Figure 2.** Photomicrograph of thin section LEW88516,6. Euhedral chromite and rounded olivine crystals poikilitically included in large orthopyroxene. Field of view is 2.2 mm.

Harvey *et al.* (1993) and used to calculate the composition of the parental melt - which was found to be similar to the calculated parental melt of other shergottites. Harvey *et al.* found that some melt inclusions contain two immiscible glasses (ratio 70:30), silica poor ~65% SiO<sub>2</sub> and extremely Si-rich (95%). Righter *et al.* (1998) determined the compositions of the melt inclusions, including the elements Rb, Ba, Mo and Ce (by ion probe).

### **Mineral Chemistry**


**Olivine:** Olivine is a major component of the poikilitic portion of LEW88516. It is compositionally zoned from

Fo<sub>70</sub> to Fo<sub>64</sub>, averaging Fo<sub>67</sub>. This is slightly more Fe-rich than the olivine in ALH77005. In both LEW88516 and ALH77005, the olivine has a distinct brown color apparently due to Fe<sup>+3</sup> produced by “shock oxidation” (Ostertag *et al.* 1984).

**Pyroxene:** Both low-Ca and high-Ca pyroxenes are present (figure 3). The large pyroxene oikiocrysts in the poikilitic portion are relatively homogeneous while the smaller pyroxenes in the non-poikilitic portion are zoned in composition. Harvey *et al.* (1993) have determined the REE and Ti, Al, Sc, Y, Zr, Cr and V contents of the different pyroxenes. Whereas the pyroxenes in LEW88516 are slightly more Fe rich, the trace element patterns are identical.

### **Mineralogical Mode**

	<b>Treiman <i>et al.</i> 1994</b>	<b>Wadhwa <i>et al.</i> 1994</b>	<b>Gleason <i>et al.</i> 1997</b>
Olivine	45.9 vol. %	50-59	57
Pyroxene	37	35	22
Maskelynite	7	8-5	16
Phosphate	0.9	1.7	<1
Chromite	0.8	0.7-2	3
Ilmenite	0.2		
Pyrrhotite	0.3		<1
Glass	7.7		



**Figure 3.** Composition diagram for pyroxene and olivine in LEW88516. Data are replotted from Harvey *et al.* (1993) and Treiman *et al.* (1994).

**Maskelynite:** Maskelynite in LEW88516 has an average composition of An<sub>52</sub> and a range from An<sub>24-58</sub> (Treiman *et al.* 1994) and is reported to contain small “bubbles”.


**Chromite:** Chromite is the most abundant accessory mineral in LEW88516. Chromite grains are commonly zoned or altered toward ulvöspinel compositions at their rims (Harvey *et al.* 1993; Gleason *et al.* 1997). Treiman *et al.* (1994) give the composition of 3 chromite grains and one grain that is a solid solution of titanomagnetite and chromite.

**Ilmenite:** Harvey *et al.* (1993), Treiman *et al.* (1994) and Gleason *et al.* (1997) give the composition of ilmenite (MgO = ~4-5 %).

**Kaersutite:** Treiman (1998), Mikouchi and Miyamoto (2000) and Monkawa *et al.* (2001) have determined the composition of “kaersutitic” amphiboles found in melt inclusions in pigeonite within LEW88516.

**Phosphate:** Whitlockite occurs as a common accessory phase in the interstices of the non-poikilitic areas of LEW88516. Harvey *et al.* (1993) determined the REE content to be about 150X C1 chondrites. The REE pattern of the whitlockite is found to be very similar to that of the bulk rock, as was also the case for ALH77005. Gleason *et al.* give the composition of whitlockite in LEW88516. Harvey *et al.* report apatite found in “melt inclusions” and give a composition.

**Sulfides:** Dreibus *et al.* (1992), Harvey *et al.* (1993) and Treiman *et al.* (1994) reported pyrrhotite in LEW88516 and Dreibus *et al.* determined that it contained 1.8 % Ni.




**Figure 4.** Normalized rare earth element diagram for LEW88516 compared with that of Shergotty. Data from Gleason *et al.* (1997).

### Whole-rock Composition

Dreibus *et al.* (1992), Treiman *et al.* (1994) and Gleason *et al.* (1997) found that the composition of LEW88516 was almost identical to that of ALH77005 (figure 4). It was found to be low, in the range of the terrestrial upper mantle (Dreibus *et al.* 1992). Analyses by Treiman *et al.* (1994) and by Warren and Kallemeyn (1996) indicate anomalous Ce. Puchel *et al.* (2009) determined the PGE and Re

### Radiogenic Isotopes

Borg *et al.* (1997, 2002) reported the crystallization age, as determined by Rb-Sr, to be  $183 \pm 710$  m.y. with an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  of  $0.71052 \pm 4$  (figure 5). Borg *et al.*



**Figure 5.** Rb-Sr isochron diagram for LEW88516 and for ALH77005. This is figure 2 in Borg *et al.* (2002).




Figure 6. Sm-Nd isochron diagram for LEW88516 and for ALH77005. This is figure 3 in Borg *et al.* (2002).

(1998, 2002) reported an Sm-Nd isochron age of  $166 \pm 16$  m.y. (figure 6). This agrees roughly with the U-Pb systematics as determined by Chen and Wasserburg (1993) who found that there was a lead-loss event  $\sim 170$  m.y.

### Cosmogenic Isotopes and Exposure Ages

Nishiizumi *et al.* (1992) and Jull *et al.* (1994) give a terrestrial exposure age of  $21 \pm 1$  thousand years.

Treiman *et al.* (1994) used the  $^{10}\text{Be}$  activity of 16.6




Figure 7. Krypton and Xenon isotopes for LEW88516 and other Martian meteorites as presented by Ott *et al.* (1996), *M&PS 31*, 103. Data for Shergotty and Zagami are open circles and data from LEW88516 are filled symbols. SPB is “shergottite parent body”.

dpm/kg from Nishiizumi *et al.* (1992) to calculate an exposure age of LEW88516 of 3.0 m.y. Schnabel *et al.* (2001) also reported  $^{10}\text{Be}$ ,  $^{26}\text{Al}$  and  $^{53}\text{Mn}$  activity. Ott and Lohr (1992) and Becker and Pepin (1993) determined  $^3\text{He}$  and  $^{21}\text{Ne}$  contents and Eugster *et al.* (1996) calculated an exposure age for LEW88516 of  $4.1 \pm 0.4$  m.y., concluding that LEW88516 was “ejected from Mars simultaneously with the other therrzolithic shergottite ALH77005 (3.4 m.y.)”

### Other Isotopes

Ott and Lohr (1992), Bogard and Garrison (1993) and Becker and Pepin (1993) determined rare gas abundances in LEW88516. Becker and Pepin (1993) and Ott *et al.* (1996) determined that the noble gasses in a glass sample from LEW88516 were in the same ratio as those in EETA79001, but much less abundant (figure 7).

Clayton and Mayeda (1996) reported the oxygen isotope composition of LEW88516 verifying that it is Martian (see figure I-3).

The initial Sr and Nd isotope ratios differ slightly from those of ALH77005 (Borg *et al.* 1997, 1998, 2002). The isotopic composition of Os has been determined by Brandon *et al.* (1997).

Mohapatra *et al.* (2009) reported the rare gas isotopes. Becker and Pepin (1993) reported that they could not find heavy nitrogen to go along with the rare gasses.

### Shock Features

Keller *et al.* (1992) and Treiman *et al.* (1994) found that the shock features in LEW88516 were very similar to those found in ALH77005. Stöffler (2000) estimated that LEW88516 was subjected to a shock pressure of  $\sim 45$  GPa and post-shock temperature of  $\sim 600$  deg. C.

### Processing

This small (13.2g) achondrite has always been processed on a laminar flow bench in the Meteorite Processing Laboratory at JSC. Most of the allocations were done a few months after classification. Some of the allocations were interior, exterior or glass-rich chips (figure 8). Most allocations were from a homogenized powder prepared from 1.66 g of interior chips by Lindstrom and Mittlefehldt. Three potted butts have been used to produce 12 thin sections.

**Table 1a. Chemical Composition of LEW88516.**

reference weight	Dreibus92	Dreibus92	Treiman94 33 mg	Treiman94 39 mg	Treiman94 19 mg	Warren96 glass-rich	Warren96	Warren96 glass-poor	Warren96								
SiO2 %	45.5	(a)				47.06											
TiO2	0.42	(a)				0.4		0.26									
Al2O3	2.99	(a)				2.36		1.23									
FeO	19.49	(a)	19.5	(a)	19.3	(a)	23	(a)	18.27								
MnO	0.47	(a)							0.48								
CaO	4.06	(a)	4.3		4.4		3.7		4.925								
MgO	25.66	(a)							23.87								
Na2O	0.49	(a)	0.536		0.558		0.7		0.386								
K2O	0.024	(a)	0.031		0.03		0.033		0.289								
P2O5		0.39	(d)														
sum	99.1																
Sc ppm	25.1	(a)	25.1	(a)	25.2	(a)	22.8	(a)	27.4	(a)	29.8	(a)					
V	180	(a)							196	(a)	198	(a)					
Cr	5672	(a)	5747	(a)	5816	(a)	5269	(a)	6900	(a)	8400	(a)					
Co	62.7	(a)	54.1	(c)	65.6	(a)	62.3	(c)	61.2	(a)	55.6	(a)					
Ni	250	(a)	280	(a)	300	(a)	300	(a)	226	(a)	244	(c)	229	(a)	230	(c)	
Cu	<80	(a)															
Zn	70	(a)	47.1	(c)			54.7	(c)	62	(a)	52	(c)	66	(a)	50	(c)	
Ga	8.4	(a)							7.6	(a)			5.8	(a)			
Ge										0.262	(c)				0.7	(c)	
As	<0.15	(a)							<0.54	(a)			<0.47	(a)			
Se	<0.7	(a)	0.22	(c)			0.35	(c)	<0.73	(a)			<1.2	(a)			
Br	0.05	(a)															
Rb		0.83	(d)	0.174	(c)		0.199	(c)									
Sr		14.7	(d)	30	(a)	20	(a)										
Y		5.69	(d)														
Zr		17.2	(d)														
Nb		0.51	(d)														
Ag ppb			4.6	(c)			7.5	(c)									
Cd ppb			9.6	(c)			15.1	(c)			<0.072	(c)			0.07	(c)	
In ppb			12.6	(c)			72.8	(c)									
Sb ppb			2.3	(c)			0.81	(c)									
Te ppb			<2	(c)			25	(c)									
I ppm	0.06	(a)															
Cs ppm		0.041	(d)	0.036	(a)	0.05	(a)	0.513	(c)								
Ba		4.93	(d)														
La	0.27	(a)	0.31	(d)	0.31	(a)	0.3	(a)	0.65	(a)	0.18	(a)		0.176	(a)		
Ce	0.94	(a)	0.87	(d)	1.4	(a)	1.9	(a)	2.4	(a)	<0.2	(a)		<0.4	(a)		
Pr																	
Nd		0.82	(d)														
Sm	0.39	(a)	0.47	(d)	0.416	(a)	0.42	(a)	0.82	(a)	0.315	(a)		0.241	(a)		
Eu	0.19	(a)	0.23	(d)	0.203	(a)	0.208	(a)	0.354	(a)	0.186	(a)		0.172	(a)		
Gd																	
Tb	0.14	(a)	0.16	(d)	0.17	(a)	0.16	(a)	0.3	(a)	0.14	(a)		0.116	(a)		
Dy	1.05	(a)	1.1	(d)													
Ho	0.19	(a)	0.24	(d)													
Er																	
Tm	0.11	(a)	0.089	(d)													
Yb	0.53	(a)	0.57	(d)	0.55	(a)	0.55	(a)	0.93	(a)	0.422	(a)		0.323	(a)		
Lu	0.078	(a)	0.083	(d)	0.076	(a)	0.076	(a)	0.14	(a)	0.071	(a)		0.059	(a)		
Hf	0.49	(a)	0.53	(d)	0.55	(a)	0.6	(a)	0.89	(a)	0.365	(a)		0.31	(a)		
Ta	0.027	(a)			30	(a)	40	(a)	<0.17	(a)				<0.17	(a)		
W ppb	<250	(a)															
Re ppb											0.071	(c)			0.064	(c)	
Os ppb											1.79	(c)			2.56	(c)	
Ir ppb	3.4	(a)		2.1	(a)	2	(a)			6.7	(a)	1.58	(c)	<5		2.32	(c)
Au ppb	0.7	(a)		0.42	(c)			0.61	(c)		0.34	(c)			0.21	(c)	
Tl ppb				4.9	(c)			4.2	(c)								
Bi ppb				1.5	(c)			3.2	(c)								
Th ppm	<0.06	(a)	0.04	(d)				0.08	(a)	<0.1	(a)			<0.13	(a)		
U ppm	0.013	(a)	0.011	(d)	0.012	(c)		0.0243	(c)								

technique: (a) INAA, (b) IDMS, (c) RNAA, (d) spark source mass spec.

**Table 1b. Composition of LEW88516 (continued).**

reference <i>weight</i>	Lodders 98 <i>average</i>	Warren 96 <i>powder</i>	Warren 96	Gleason 97 <i>147 mg</i>	Gleason 97 <i>147 mg.</i>	Brandon 2000 <i>103.8 mg.</i>	Borg 2001 <i>25 mg.</i>
SiO <sub>2</sub>	46						
TiO <sub>2</sub>	0.39	0.32		0.36	(a) 0.45	(a)	
Al <sub>2</sub> O <sub>3</sub>	3.31	2.93		3.45	(a) 3.86	(a)	
FeO	19	18.91		20.9	(a) 20.3	(a)	
MnO	0.49	0.48		0.51	(a) 0.51	(a)	
CaO	4.2	4.114		4.25	(a) 4.54	(a)	
MgO	25	25.5		23.7	(a) 24.9	(a)	
Na <sub>2</sub> O	0.56	0.516		0.588	(a) 0.595	(a)	
K <sub>2</sub> O	0.029	0.301		0.028	(a) 0.028	(a)	
P <sub>2</sub> O <sub>5</sub>	0.39						
sum							
Sc ppm	25	25.2	(a)	26.7	(a) 26.5	(a)	
V	170	143	(a)	171	(a) 178	(a)	
Cr	5880	5800	(a)	6295	(a) 6568	(a)	
Co	65	66.7	(a)	66.5	(a) 64.1	(a)	
Ni	280	245	(a) 254	(c) 315	(a) 276	(a)	
Cu							
Zn	60	69	(a) 54	(c) 62.1	(a) 60.6	(a)	
Ga	9.2	8.7	(a)	9.1	(a) 10.4	(a)	
Ge	0.6		0.6	(c)			
As	<0.93	<0.93	(a)				
Se	0.34	<0.64	(a)	0.35	(a) 0.44	(a)	
Br	0.05						
Rb	1.1			1.41	(a) 0.97	(a)	0.657 (a)
Sr	17			14.3	(a) 17.7	(a)	12.9 (a)
Y	5.7						
Zr	13			12.4	(a) 9.6	(a)	
Nb	0.51						
Mo							
Pd ppb							
Ag ppb	6.1						
Cd ppb	12.4		<0.08				
In ppb	12.6 or 72.8						
Sb ppb	1.6						
Te ppb	25						
I ppm	0.06						
Cs ppm	0.055			0.079	(a) 0.06	(a)	
Ba	4.9						
La	0.33	0.3	(a)	0.41	(a) 0.41	(a)	
Ce	1.26	2.1	(a)	0.8	(a) 0.82	(a)	
Pr							
Nd	0.82						0.447 (b)
Sm	0.44	0.4	(a)	0.474	(a) 0.502	(a)	0.267 (b)
Eu	0.22	0.221	(a)	0.242	(a) 0.256	(a)	
Gd							
Tb	0.16	0.14	(a)	0.167	(a) 0.172	(a)	
Dy	1.08						
Ho	0.23			0.247	(a) 0.23	(a)	
Er							
Tm	0.1						
Yb	0.57	0.5	(a)	0.658	(a) 0.656	(a)	
Lu	0.083	0.079	(a)	0.091	(a) 0.097	(a)	
Hf	0.52	0.43	(a)	0.514	(a) 0.548	(a)	
Ta	0.035	<0.22	(a)		0.041	(a)	
W ppb	140			0.11	(a) 0.17	(a)	
Re ppb	<0.09		0.09 (c)				0.0222 (b)
Os ppb	3.3		3.3 (c)				0.905 (b)
Ir ppb	3		3.1 (c)	3.6	(a) 3.8	(a)	
Au ppb	0.53		0.39 (c)				
Tl ppb	4.6						
Bi ppb	2.4						
Th ppm	0.037	<0.09	(a)		0.044	(a)	
U ppm	0.014						

technique: (a) INAA, (b) IDMS, (c) RNAA




Figure 8. Genealogy diagram for LEW88516 for samples allocated up to 1997.

LEW88516 is listed as a “restricted” sample by the MWG (Score and Lindstrom 1993, page 5), because of its small size.

#### References for LEW88516

- Ariskin A.A. (1997) Parent magmas of SNC harzburgites: Phase equilibria modeling (abs). *Lunar Planet. Sci. XXVIII*, 51-52. Lunar Planetary Institute, Houston.
- Becker R.H. and Pepin R.O. (1993a) Nitrogen and Noble gases in a glass sample from LEW88516 (abs). *Lunar Planet. Sci. XXIV*, 77-78. Lunar Planetary Institute, Houston.
- Becker R.H. and Pepin R.O. (1993b) Nitrogen and Noble gases in a glass sample from the LEW88516 shergottite (abs). *Meteoritics* **28**, 637-640.
- Bogard D.D. and Garrison D.H. (1993) Noble gases in LEW88516 Shergottite: Evidence for exposure age pairing with ALHA77005 (abs). *Lunar Planet. Sci. XXIV*, 139-140. Lunar Planetary Institute, Houston.
- Borg L.E., Nyquist L.E. and Wiesmann Henry (1998a) Rb-Sr isotopic systematics of the Lherzolithic Shergottite LEW88516 (abs#1233). *Lunar Planet. Sci. XXIX* Lunar Planetary Institute, Houston.
- Borg L.E., Nyquist L.E., Wiesmann H. and Reese Young (1998b) Samarium-neodymium isotopic systematics of the Lherzolithic Shergottite Lewis Cliff 88516 (abs). *Meteorit. & Planet. Sci.* **33**, A20.
- Borg L.E., Nyquist L.E., Wiesmann H. and Reese Y. (2001b) Rubidium-strontium and Samarium-neodymium isotopic systematics of the Lherzolithic Shergottites ALH77005 and LEW88516: Constraints on the petrogenesis of Martian meteorites (abs). *Meteorit. & Planet. Sci.* **36**, A25.
- Borg L.E., Nyquist L.E., Wiesmann H. and Reese Y. (2002a) Constraints on the petrogenesis of Martian meteorites from the Rb-Sr and Sm-Nd isotopic systematics of the Lherzolithic Shergottites ALH77005 and LEW88516. *Geochim. Cosmochim. Acta* **66**, 2037-2053.
- Boynton W.V., Hill D.H. and Kring D.A. (1992) The trace-element composition of LEW88516 and its relationship to SNC meteorites (abs). *Lunar Planet. Sci. XXIII*, 147-148. Lunar Planetary Institute, Houston.
- Brandon A.D., Walker R.J., Morgan J.W. and Goles G.G. (2000a) Re-Os isotopic evidence for early differentiation of the Martian mantle (abs#1676). *Lunar Planet. Sci. XXXI*, Lunar Planetary Institute, Houston.
- Brandon A.D., Walker R.J., Morgan J.W. and Goles G.G. (2000b) Re-Os isotopic evidence for early differentiation of the Martian mantle. *Geochim. Cosmochim. Acta* **64**, 4083-4095.
- Chen J.H. and Wasserburg G.J. (1993) LEW88516 and SNC meteorites (abs). *Lunar Planet. Sci. XXIV*, 275-276. Lunar

- Planetary Institute, Houston. of Lherzolite Yamato-793605. *Antarctic Meteorite Research* **10**, 141-149. Nat. Inst. Polar Res., Tokyo.
- Clayton R.N. and Mayeda T.K. (1996) Oxygen isotopic studies of achondrites. *Geochim. Cosmochim. Acta* **60**, 1 9 9 9 - 2 0 1 7 .
- Delaney J.S. (1992) Petrological comparison of LEW88516 and ALHA77005 Shergottites (abs). *Meteoritics* **27**, 213-214.
- Delaney J.S., Sutton S. R. and Dyar D.M. (1998) Variable oxidation states of iron in Martian meteorites (abs#1241). *Lunar Planet. Sci.* **XXIX** Lunar Planetary Institute, Houston .
- Delaney J.S. and Dyar M.D. (2000) Correction of the calibration of ferric/ferrous determinations in pyroxene from Martian samples and achondritic meteorites by synchrotron microXANES spectroscopy. (abs). *Lunar Planet. Sci.* **XXXI**, Lunar Planetary Institute, Houston.
- Douglas C., Wright I.P., Yates P.D. and Pillinger C.T. (1992) The carbon isotopic composition of LEW88516, the fifth shergottite (abs). *Meteoritics* **27**, 215-216.
- Dreibus G., Jochum K.H., Palme H., Spettel B., Wlotzka F. and Wänke H. (1992) LEW88516: A meteorite compositionally close to the "Martian mantle" (abs). *Meteoritics* **27**, 216-217.
- Dreibus G., Wänke H. and Lugmair G.W. (1997) Volatile inventories of Mars and Earth and their implications for the evolution of the planetary atmospheres (abs). In *Conference on Early Mars: Geologic and hydrologic evolution, physical and chemical environments, and the implications for life.* (eds. Clifford et al.) *LPI Contribution* **916**, 26. Lunar Planetary Institute, Houston.
- Eugster O. and Weigel A. (1992) Exposure histories of lodranites, Shergottite LEW88516 and CK-chondrites (abs). *Meteoritics* **27**, 219.
- Eugster O., Weigel A. and Polnau E. (1996) Two different ejection events for basaltic Shergottites QUE94201, Zagami and Shergotty (2.6 Ma ago) and Lherzolic Shergottites LEW88516 and ALHA77005 (3.5 Ma ago) (abs). *Lunar Planet. Sci.* **XXVII**, 345-346. Lunar Planetary Institute, Houston.
- Eugster O. and Polnau E. (1997a) Mars-Earth transfer time of Lherzolite Yamato-793605 (abs). *Proc. NIPR Sym. Antarctic Meteorites* **22nd**, 31-33. Nat. Inst. Polar Res., Tokyo.
- Eugster O. and Polnau E. (1997b) Mars-Earth transfer time
- Eugster O., Weigel A. and Polnau E. (1997a) Ejection times of Martian meteorites. *Geochim. Cosmochim. Acta* **61**, 2749-2757.
- Gleason J.D., Kring D.A., Hill D.H. and Boynton W.V. (1997c) Petrography and bulk chemistry of Martian Lherzolite LEW88516. *Geochim. Cosmochim. Acta* **61**, 4007-4014.
- Goodrich C.A. and Harvey R.P. (2002) The parent magmas of Lherzolic Shergottites ALH77005 and LEW88516: A reevaluation from magmatic inclusions in olivine and chromite (abs). *Meteorit. & Planet. Sci.* **37**, A54.
- Greenwood J.P., Riciputi L.R. and McSween H.Y. (1997b) Sulfide isotopic compositions in Shergottites and ALH84001 and possible implications for life on Mars. *Geochim. Cosmochim. Acta* **61**, 4449-4453.
- Grossman J.N. (1994) The Meteoritical Bulletin, No. 76: The U. S. Antarctic meteorite collection. *Meteoritics* **29**, 1 0 0 - 1 4 3 .
- Harvey R.P. and McSween H.Y. (1992a) The mineralogy and petrography of LEW88516 (abs). *Meteoritics* **27**, 231-2 3 2 .
- Harvey R.P., Wadhwa M., McSween H.Y. and Crozaz G. (1993) Petrography, mineral chemistry and petrogenesis of Antarctic Shergottite LEW88516. *Geochim. Cosmochim. Acta* **57**, 4769-4783.
- Herd C.D.K., Jones J.H., Schearer C.K. and Papike J.J. (2001a) Systematics of Ni, Co, Cr and V in olivine from planetary melt systems: Martian basalts (abs#1635). *Lunar Planet. Sci.* **XXXII** Lunar Planetary Institute, Houston.
- Herd C.D.K., Karner J.M., Shearer C.K. and Papike J.J. (2001c) The effect of oxygen fugacity on Co and Ni partitioning in olivine: Insights into Martian magmas (abs). *Meteorit. & Planet. Sci.* **36**, A78-79.
- Jull A.J.T., Cielaszyk E., Brown S.T. and Donahue D.J. (1994b) <sup>14</sup>C terrestrial ages of achondrites from Victoria Land, Antarctica (abs). *Lunar Planet. Sci.* **XXV**, 647-648. Lunar Planetary Institute, Houston.
- Keller L.P., Treiman A.H. and Wentworth S.J. (1992) Shock effects in the Shergottite LEW88516: Optical and electron microscope observations (abs). *Meteoritics* **27**, 242.
- Kring D.A. and Gleason J.D. (1999) Silicous igneous rock on Mars (abs#1611). *Lunar Planet. Sci.* **XXX**, Lunar Planetary Institute, Houston.

- Kurihara T., Mikouchi T., Saruwatari K., Kameda J., Ari T., Hoffmann V. and Miyamoto M. (2008a) Transmission electron microscopy of “brown” color olivines in Martian and lunar meteorites (abs#2478). *Lunar Planet. Sci.* **XXXIX** Lunar Planetary Institute, Houston.
- Kurihara T., Mikouchi T., Saruwatari K., Kameda J. and Miyamoto M. (2008b) Transmission electron microscopy of olivine in the LAR06319 olivine-phyrlic Shergottite (abs#5177). *Meteorit. & Planet. Sci.* **43**, A83.
- Lindstrom M.M., Mittlefehldt D.W., Treiman A.H., Wentworth S.J., Gooding J.L., Keller L.P. and McKay G. A. (1992) LEW88516: A new Shergottite from Antarctica (abs). *Lunar Planet Sci.* **XXIII**, 783-784. Lunar Planetary Institute, Houston.
- Mikouchi T. and Miyamoto M. (1996b) Comparative mineralogy of Antarctic Lherzolitic Shergottites Allan Hills 77005, Lewis Cliff 88516 and Yamoto 793605 (abs). *Meteorit. & Planet. Sci.* **31**, A89-A90.
- Mikouchi T., Miyamoto M. and McKay G. A. (1997a) Similarities in zoning of pyroxenes from QUE94201 and EETA79001 Martian meteorites (abs). *Lunar Planet. Sci.* **XXVIII**, 955-956. Lunar Planetary Institute, Houston.
- Mikouchi T. and Miyamoto M. (1999b) Micro Raman spectroscopy of amphiboles and Al-Ti-rich pyroxenes in the Martian meteorites Zagami and LEW88516 (abs#1559). *Lunar Planet. Sci.* **XXX** Lunar Planetary Institute, Houston.
- Mikouchi T. and Miyamoto M. (2000a) Lherzolitic Martian meteorites Allan Hills 77005, Lewis Cliff 88516 and Yamoto-793605: Major and minor element zoning in pyroxene and plagioclase glass. *Antarct. Meteorite Res.* **13**, 256-269.
- Mikouchi T. and Miyamoto M. (2000b) Micro Raman spectroscopy of amphiboles and pyroxenes in the Martian meteorites Zagami and Lewis Cliff 88516. *Meteorit. & Planet. Sci.* **35**, 155-159.
- Mittlefehldt D.W., Wentworth S.J., Wang M.-S., Lindstrom M.M. and Lipschutz M.E. (1997b) Geochemistry of and alteration phases in Martian Lherzolite Y-793605. *Antarctic Meteorite Res.* **10**, 109-124. Nat. Inst. Polar Res., Tokyo.
- Mohapatra R.K., Schwenzer S.P., Herrmann S., Murty S.V.S., Ott U. and Gilmour J.D. (2009) Noble gases and nitrogen in Martian meteorites Dar al Gani 476, Sayh al Uhaymir 005 and Lewis Cliff 88516: EFA and extra neon. *Geochim. Cosmochim. Acta* **42**, 131-148.
- Monkawa A., Makino K., Ishii T., Ohtsuki M., Mikouchi T. and Miyamoto M. (2001) The determination of Fe<sup>2+</sup>/Fe ratios of kaersutite in the Martian meteorite LEW88516 by electron microprobe (abs). *Meteorit. & Planet. Sci.* **36**, A140.
- Monkawa A., Mikouchi T., Miyamoto M., Koisumi E., Miyata Y. and Ohsumi K. (2002c) On the formation of Ti-rich kaersutite amphiboles in Martian meteorites (abs). *Antarctic Meteorites XXVII*, 102-104. Nat. Inst. Polar Res., Tokyo
- Nishiizumi K., Arnold J.R., Caffee M.W., Finkel R.C. and Southon J. (1992) Exposure histories of Calalong Creek and LEW88516 meteorites (abs). *Meteoritics* **27**, 270.
- Ott U. and Löhr H.-P. (1992) Noble gases in the new (now old) Shergottite LEW88516 (abs). *Meteoritics* **27**, 271.
- Ott U., Löhr H.-P. and Begemann F. (1996) Etching and crushing SNCs: More Noble gas data (abs). *Meteorit. & Planet. Sci.* **31**, A103.
- Papike J.J., Karner J.M., Spilde M.N., Shearer C.K. and Burger P.V. (2009b) Silicate mineralogy of Martian meteorites. *Geochim. Cosmochim. Acta* **73**, 7443-7485. (invited review with great pictures of textures)
- Puchtel I.S., Walker R.J., Brandon A.D. and Irving A.J. (2008) Highly siderophile element abundances in SNC meteorites: An update (abs#1650). *Lunar Planet. Sci.* **XXXIX** Lunar Planetary Institute, Houston .
- Righter K., Hervig R.L. and Kring D.A. (1997) Ion microprobe analyses of SNC meteorite melt inclusions (abs). *Lunar Planet. Sci.* **XXVIII**, 1181-1182. Lunar Planetary Institute, Houston.
- Righter K., Hervig R.L. and Kring D.A. (1998) Accretion and core formation on Mars: Molybdenum contents of melt inclusion glasses in three SNC meteorites. *Geochim. Cosmochim. Acta* **62**, 2167-2177.
- Satterwhite C. and Mason B. (1991) Macroscopic and thin section description of LEW88516. In *Antarctic Meteorite Newsletter* **14** (2), 19. JSC Curator's Office, Houston.
- Schnabel C., Ma. P., Herzog G.F., Faestermann T., Knie K. and Korschinek G. (2001) <sup>10</sup>Be, <sup>26</sup>Al and <sup>53</sup>Mn in Martian meteorites (abs#1353). *Lunar Planet. Sci.* **XXXII**, Lunar Planetary Institute, Houston.
- Shearer C.K., Burger P.V., Papike J.J. and Karner J. (2009) Comparison between RBT04262 and Lherzolitic Shergottites (ALHA77005 and LEW88516) (abs#1300).

*Lunar Planet. Sci.* **XL**, Lunar Planetary Institute, The Woodlands.

Treiman A.H. (1993c) Xenoliths in the EETA79001 Shergottite: Geological and astronomical implications of similarities to the ALHA77005 and LEW88516 Shergottites (abs). *Meteoritics* **28**, 451.

Treiman A.H. (1995a)  $S \neq NC$ : Multiple source areas for Martian meteorites. *J. Geophys. Res.* **100**, 5329-5340.

Treiman A.H. (1998b) Amphiboles in more Martian meteorites: Elephant Moraine 79001B, Elephant Moraine 79001X and Lewis Cliff 88516 (abs). *Meteorit. & Planet. Sci.* **33**, A156.

Treiman A.H., McKay G.A., Bogard D.D., Wang M.-S., Lipschutz M.E., Mittlefehldt D.W., Keller L., Lindstrom M.M. and Garrison D. (1994a) Comparison of the LEW88516 and ALHA77005 Martian meteorites: Similar but distinct. *Meteoritics* **29**, 581-592.

Wadhwa M. and Crozaz G. (1992c) Trace element characteristics of the Shergottite LEW88516 (abs). *Meteoritics* **27**, 302-303.

Wadhwa M. and Crozaz G. (1993) Rare earth elements in individual minerals in Shergottites (abs). *Lunar Planet. Sci.* **XXIV**, 1473-1474. Lunar Planetary Institute, Houston.

Wang M.-S., Mokoš J. and Lipschutz M.E. (1997) Volatile and other trace elements in Martian meteorites (abs). *Lunar Planet. Sci.* **XXVIII**, 1493-1494. Lunar Planetary Institute, Houston.

Wentworth S.J. and Gooding J.L. (1993) Weathering features and secondary minerals in Antarctic Shergottites ALHA77005 and LEW88516 (abs). *Lunar Planet. Sci.* **XXIV**, 1507-1508. Lunar Planetary Institute, Houston.

Wright I.P., Douglas C. and Pillinger C.T. (1993a) The carbon components in SNC meteorites of feldspathic harzburgite composition (abs). *Lunar Planet. Sci.* **XXIV**, 1539-1540. Lunar Planetary Institute, Houston.

Wright I.P., Douglas C. and Pillinger C.T. (1993b) Further carbon isotope measurements of LEW88516 (abs). *Lunar Planet. Sci.* **XXIV**, 1541-1542. Lunar Planetary Institute, Houston.