


**76275**  
Impact Melt Breccia  
55.9 grams



*Figure 1: Side view of 76275. Cube is 1 cm. S73-15077*



*Figure 2: Original location of 76275 on side of block 1, of boulder at station 6, Apollo 17. AS17-140-21443.*



Figure 3: Photomicrograph of thin section 76275,4 by C Meyer. Partially crossed polarizers. Field of view about 2 mm.

### **Introduction**

76275 was chipped off of the side of block 1 of the big boulder at station 6, Apollo 17 (Wolfe et al. 1981). One side is freshly broken, the other side rounded by micrometeorite bombardment and coated with patina (glass splash from the regolith).

The remaining large piece of 76275 contains white clasts that have never been studied.

### **Petrography**

76275 is a clast-bearing impact melt breccia (figure 3), similar to other samples of the station 6 boulder (Phinney 1981). The matrix has a poikilitic texture with a matrix of interlocking low-Ca pyroxene oikocrysts that surround abundant clasts and laths of plagioclase, olivine and augite. Sieve-like ilmenite is found between pyroxene oikocrysts.

The composition of pyroxene and olivine is given in figure 5. Misra et al. (1976) studied iron grains in 76275 (figure 6).

### **Chemistry**

The bulk composition of 76275 was reported by Phinney (1981). Gros et al. (1976) reported the critical siderophile elements, showing that 76275 was similar to the rest of the station six boulder.

### **Radiogenic age dating**

Cadogen and Turner (1976) determined an age of  $4.02 \pm 0.04$  b.y. by the Ar/Ar plateau technique (figure 7).

### **Cosmogenic isotopes and exposure ages**

Rancitelli et al. (1974) determined the cosmic-ray-induced activity of  $^{22}\text{Na} = 100$  dpm/kg,  $^{26}\text{Al} = 110$  dpm/




Figure 4: Photomicrographs of 76275,46. PPL-S79-27774, XPL-75, RL-73. Field of view is 1.4 mm.

kg,  $^{46}\text{Sc} = 7 \text{ dpm/kg}$ ,  $^{54}\text{Mn} = 103 \text{ dpm/kg}$  and  $^{56}\text{Co} = 86 \text{ dpm/kg}$ .

### Processing

76275 was to be part of a consortium study by Bill Phinney and team. However, the processing was delayed and, for whatever reason, the sample studies were not completed. There are 16 thin sections of 76275. It has been sawn to create an end piece and a wedge which were further subdivided (figure 8).




Figure 5: Pyroxene and olivine in 76275 as reported by Phinney (1981).




Figure 6: Ni and Co content of metal grains in 76275 (Misra et al. 1976).




Figure 7: Ar/Ar plateau diagram for Station 6 samples (Cadogen and Turner 1976)

### Summary of Age Data for 76275

Ar/Ar  
Cadogen and Turner 1976  $4.05 \pm 0.04 \text{ b.y}$

**Table 1. Chemical composition of 76275.**

reference weight	Gros76	Rancitelli74	Simonds81	tan matrix
SiO2 %			47.14	(c)
TiO2			1.65	(c)
Al2O3			18.7	(c)
FeO			8.54	(c)
MnO				
MgO			9.22	(c)
CaO			12.06	(c)
Na2O			0.72	(c)
K2O		0.27	(b) 0.34	(c)
P2O5				
S %				
sum				
Sc ppm				
V				
Cr				
Co				
Ni	387	(a)		
Cu				
Zn	4	(a)		
Ga				
Ge ppb	383	(a)		
As				
Se	125	(a)		
Rb	3.67	(a)		
Sr				
Y				
Zr				
Nb				
Mo				
Ru				
Rh				
Pd ppb	19.8	(a)		
Ag ppb	1.22	(a)		
Cd ppb	8.8	(a)		
In ppb	12.4	(a)		
Sn ppb				
Sb ppb	2	(a)		
Te ppb	9.8	(a)		
Cs ppm	0.196	(a)		
Ba				
La				
Ce				
Pr				
Nd				
Sm				
Eu				
Gd				
Tb				
Dy				
Ho				
Er				
Tm				
Yb				
Lu				
Hf				
Ta				
W ppb				
Re ppb	0.725	(a)		
Os ppb	8.6	(a)		
Ir ppb	7.76	(a)		
Pt ppb				
Au ppb	5.1	(a)		
Th ppm			5.69	(b)
U ppm	2.35	(a)	1.4	(b)

technique: (a) RNAA, (b) rad. count. , (c) unreliable fused bead e. probe





Figure 8a and b: Processing photo of 76275. Cube is 1 cm. S75-24192 and S75-24194



### References for 76275

Butler P. (1973) Lunar Sample Information Catalog Apollo 17. Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.

Cadogan P.H. and Turner G. (1976) The chronology of the Apollo 17 Station 6 boulder. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2267-2285.

Gros J., Takahashi H., Hertogen J., Morgan J.W. and Anders E. (1976) Composition of the projectiles that bombarded the lunar highlands. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2403-2425.

Meyer C. (1994) **Catalog of Apollo 17 rocks**: Volume 4. Curator's Office JSC 26088 pp. 644

Misra K.C., Walker B.M. and Taylor L.A. (1976) Textures and compositions of metal particles in Apollo 17, Station 6 boulder samples. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2251-2266.

Phinney W.C., Consortium Leader (1974) Progress report: Apollo 17, station 6 boulder consortium (abs). *Lunar Sci.* V, Suppl. A. The Lunar Science Institute, Houston.

Rancitelli L.A., Perkins R.W., Felix W.D. and Wogman N.A. (1974) Solar flare and lunar surface process characterization at the Apollo 17 site. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2185-2203.

Simonds C.H. (1975) Thermal regimes in impact melts and the petrology of the Apollo 17 Station 6 boulder. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 641-672.

Simonds C.H., Warner J.L. Phinney W.C. and McGee P.E. (1976) Thermal model for impact breccia lithification: Manicouagan and the moon. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2509-2528.

Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L. and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.